

# Research Statement

Thotsaporn Aek Thanatipanonda

August 3, 2013

My research interest is in automated proving of theorems in combinatorics. Or at least using computers to do their best assisting humans to solve the problems. This includes: making conjectures, giving numerical evidence, and doing symbolic computations. I like many areas of combinatorics, my favorites being Combinatorial Game theory, Ramsey theory, and the theory of partitions.

## 1 Dissertation Research

Richard Hamming, a great applied mathematician of the last century, said “the purpose of computing is insight, not numbers”. Understanding makes me feel good. Computation gives me insight in two different ways. First the *act* of programming makes me understand the problem much better, and at a deeper level, and second, the *output* often leads to further understanding. Once we collect all the information, we see the *big picture* without worrying about the details of calculations. I like to solve challenging problems, and it is always the case that knowing computer programming helps the calculational part goes smoother.

During my graduate years, I learned the philosophy and methodology of using computers in mathematical research, specifically symbolic computation and automated proving, from Prof. Doron Zeilberger, my advisor. In my opinion, it is not very important in what mathematical area I am working on, since the *method* and *experience* gained in doing computer-assisted and computer-generated research in one area are likely to be transferable to other areas.

**The “Toads and Frogs” game:** I do research in combinatorial game theory. The modern theory was developed by J.Conway, E.Berlekamp, R.Guy, who wrote the classic book *Winning Ways*, that mostly dealt with *partizan* games, and by Aviezri Fraenkel and his many students, who study *impartial* games.

I started my research by investigating “Toads and Frogs”. In 1994, Jeff Erickson made five intriguing conjectures [3]. In my paper (joint with my advisor), [15], we proved one of Erickson’s conjectures. In another paper of mine, [14], [12], I settled three others, two positively and one

negatively. In the process, I proved much more general results. More importantly, we discovered a new technique that allowed computers to conjecture and prove values of many families of positions, completely automatically. We called it the “Finite State Method”. In the past, mathematicians had a hard time proving the values of positions by logical reasoning. Once computers are taught how to do this “reasoning”, they can do much more, of course. I implemented all this in the computer algebra system Maple. My program can also determine a winning move (if it exists) in each position, by using the recurrence relation for each of the positions.

**Problems on Schur Triples:** Schur’s theorem is one of the “super six” theorems in Ramsey theory mentioned in Ron Graham, Bruce Rothschild, and Joel Spencer’s famous monograph. The *Schur number*,  $s(r)$ , is the smallest number such that for any  $r$ -coloring of the set  $1, 2, \dots, s(r)$ , there must be a monochromatic solution to the equation  $x + y = z$ . A triple  $(x, y, z)$  is called a *Schur triple*. Only the first four values are currently known:  $s(1) = 2, s(2) = 5, s(3) = 14, s(4) = 45$ . Even  $s(5)$  is unknown. It is a natural question, asked by Ron Graham, what is the least number of Schur triples in  $\{1, \dots, n\}$  in a 2-coloring, as  $n \rightarrow \infty$ . This problem was solved independently by [7], [8], who shared the 100\$ prize offered by Graham. Yet another proof appeared later [2]. The answer is  $\frac{n^2}{22} + O(n)$ .

Ron Graham further asked about the minimum number of monochromatic triples  $(x, y, x + ay)$  in 2-colorings of  $[1, n]$ . I wrote a Maple program to find the optimal coloring when  $n$  is small. After finding the pattern, I developed a method that I called “greedy calculus”, to calculate “good” colorings which gives upper bounds. Then I used another method involving calculus that gives lower bounds. So far the upper bounds and lower bounds do not match. But these upper bounds are conjectured to be sharp. I also applied this algorithm to the original Schur triples, but with  $r$ -colorings, to get upper bounds which I suspect to be sharp. These considerably improve previous bounds due to A. Robertson. These results are in [13].

**Moment Calculus:** The expectation functional is a powerful tool to study combinatorial objects, and often gives you quite useful information. To find higher moments, the computation gets complicated and we need computers to do symbolic computation for us. The technique has already been demonstrated in [16], [17], [18], [1]. Once we find high enough moments, they could actually be useful for calculating lower bounds for enumerating combinatorial objects (see [17]). I have calculated the higher moments of the Ramsey graph of  $K_3$  and  $K_4$  on  $n$  vertices for the  $r$ -coloring of edges of the complete graph, the second moment of Schur Triples with  $r$ -coloring on  $[1, n]$ . The results are on my website.

## 2 Post Graduate School

**Evaluating Determinant:** In the mid 1970’s, Mathematician started to relate counting problems from combinatorics, especially in the area of plane partitions, to the evaluation of certain determinant. During the Montreal Conference in 1985, Richard Stanley raised the problems concerned the enumeration of “symmetry classes” of plane partitions which later be published, [9] and [10].

Today all of the problems in [10] were solved. The last standing problems was solved earlier this year in [6] using Zeilberger Algorithm. I have a chance to work with Koutchen, one of the author of [6], in which a collaboration turned out to be fruitful. We solved three six-years-old-conjectures published by Christian Krattenthaler in his famous article, [5], related to evaluating determinants. The results are in [4].

**Ramsey Multiplicity:** As a problem along the same lines as my work on (generalized) Schur triples mentioned above, we can ask a similar problem related to Ramsey numbers instead of Schur numbers. For a fixed number  $k$ , find the least number of monochromatic  $K_k$  for any 2-edge-coloring graph in  $K_n$  where  $n \rightarrow \infty$ . The easy solution for  $K_3$  is known. The answer is  $\frac{n^3}{24} + O(n^2)$  which is the same as the average. The answer for  $k \geq 4$  is unknown. I observed some general results of an  $r$ -edge-coloring graph where  $r \geq 2$  in [11].

### 3 Future Goals

Using computers in mathematical research is getting more and more common, but most of it is done in “interactive” mode. I believe that a more systematic and methodical approach, with extensive programming, can lead to even more.

I want to keep working in the kind of research compatible with my strengths. Experimental mathematics and symbolic computation are the *key words*. I will try and focus on problems where there is a way for computers to take part in formulating conjectures, delivering numerical results for small examples, doing symbolic computation, and whenever possible, proving interesting new results.

I am willing to work on anything, and like to collaborate also.

### References

- [1] Andrew Baxter and Doron Zeilberger. The number of inversions and the major index of permutations are asymptotically joint-independently normal (second edition!). *Personal Journal of Shalosh B. Ekhad and Doron Zeilberger*, 2010.
- [2] Boris A. Datskovsky. On the number of monochromatic Schur triples. *Adv. in Appl. Math.*, 31(1):193–198.
- [3] Jeff Erickson. New toads and frogs results. *Games of no chance (Berkeley, CA, 1994)*, volume 29 of Math. Sci. Res. Inst. Publ.:299–310, 1996.
- [4] Christoph Koutschan and Thotsaporn “Aek” Thanatipanonda. Advanced computer algebra for determinants. *preprint (available from the author’s website)*.

- [5] Christian Krattenthaler. Advanced determinant calculus: A complement. *Linear Algebra and its Applications*, 411:68–166, 2005.
- [6] Christoph Koutschan Manuel Kauers and Doron Zeilberger. A proof of george andrews’ and david robbins’ q-tspp conjecture. *Proceedings of the National Academy of Science*.
- [7] Aaron Robertson and Doron Zeilberger. A 2-coloring of  $[1, n]$  can have  $(1/22)n^2 + O(n)$  monochromatic Schur triples, but not less! *Electron. J. Combin.*, 5:Research Paper 19, 4pp. (electronic), 1998.
- [8] Tomasz Schoen. The number of monochromatic Schur triples. *European J. Combin.*, 20(8):855–866.
- [9] Richard P. Stanley. A baker’s dozen of conjectures concerning plane partitions. *Combinatoire Enumerative*, Lecture Notes in Math., no. 1234:285–293, 1986.
- [10] Richard P. Stanley. Symmetries of plane partitions. *J. Combinatorial Theory (A)*, 43:103–113, 1986.
- [11] Thotsaporn “Aek” Thanatipanonda. On the ramsey multiplicity of complete graphs. *Personal Note (available from the author’s website)*.
- [12] Thotsaporn “Aek” Thanatipanonda. Three results in combinatorial game Toads and Frogs. *preprint (available from the author’s website)*.
- [13] Thotsaporn “Aek” Thanatipanonda. On the monochromatic Schur triples type problem. *Electron. J. Combin*, 16(1):R(14), 2009.
- [14] Thotsaporn “Aek” Thanatipanonda. Further hopping with Toads and Frogs. *Electronic Journal of Combinatorics*, 18(1):R(14), 2011.
- [15] Thotsaporn “Aek” Thanatipanonda and Doron Zeilberger. A symbolic finite-state approach for automated proving of theorems in combinatorial game. *J. of Difference Equations and Applications*, 15:111–118, 2009.
- [16] Doron Zeilberger. Symbolic moment calculus. I. Foundations and permutation pattern statistics. *Ann. Comb.*, 8(3):369–378, 2004.
- [17] Doron Zeilberger. Symbolic moment calculus. II. Why is ramsey theory sooooo eeenormaously hard? *Combinatorial Number Theory*, 2007.
- [18] Doron Zeilberger. The automatic central limit theorems generator (and much more!). *Advances in Combinatorial Mathematics: Proceedings of the Waterloo Workshop in Computer Algebra 2008 in honor of Georgy P. Egorychev*, 2009.